

Higher Genus Correlators from the Hermitian One-Matrix Model

J. Ambjørn¹

*The Niels Bohr Institute
Blegdamsvej 17, DK-2100 Copenhagen, Denmark*

L. Chekhov²

*Steklov Mathematical Institute
Vavilov st.42, GSP-1, 117966 Moscow, Russia*

Yu. Makeenko³

*Institute of Theoretical and Experimental Physics
B.Cheremuskinskaya 25, 117259 Moscow, Russia*

Abstract

We develop an iterative algorithm for the genus expansion of the hermitian $N \times N$ one-matrix model (= the Penner model in an external field). By introducing moments of the external field, we prove that the genus g contribution to the m -loop correlator depends only on $3g - 2 + m$ lower moments ($3g - 2$ for the partition function). We present the explicit results for the partition function and the one-loop correlator in genus one. We compare the correlators for the hermitian one-matrix model with those at zero momenta for $c = 1$ CFT and show an agreement of the one-loop correlators for genus zero.

Submitted to *Physics Letters B*

¹E-mail: ambjorn@nbivax.nbi.dk

²E-mail: chekhov@qft.mian.su

³E-mail: makeenko@nbivax.nbi.dk / makeenko@desyvax.bitnet / makeenko@itep.msk.su

1 Introduction

Recently there has renewed interest in the hermitian one-matrix model which is defined by the partition function

$$Z_N[t.] = \int DX \, e^{-\sum_{k=0}^{\infty} t_k \operatorname{tr} X^k} \quad (1.1)$$

where the integration goes over $N \times N$ hermitian matrices. As is proven in Ref.[1], this model is equivalent as $N \rightarrow \infty$ to the following model in an external field:

$$Z_N[\eta; \alpha] = e^{-\frac{N}{2} \operatorname{tr} \eta^2} \int DX \, e^{N \operatorname{tr} \left(-\frac{1}{2} X^2 + \eta X + \alpha \log X \right)}, \quad (1.2)$$

with η and $\{t_k\}$ being related by the Miwa transformation

$$t_k = \frac{1}{k} \operatorname{tr} \eta^{-k} - \frac{N}{2} \delta_{k2} \quad \text{for } k \geq 1, \quad t_0 = \operatorname{tr} \log \eta^{-1}. \quad (1.3)$$

The partition function (1.2), in turn, is associated [2] with an external field problem for the Penner model [3]:

$$Z_N[\Lambda; \alpha] = e^{-\frac{\alpha^2 N}{2} \operatorname{tr} \Lambda^{-2}} (\det \Lambda)^{N(\alpha+1)} \int DX \, e^{N \operatorname{tr} \left(-\frac{1}{2} \Lambda X \Lambda X + \alpha [\log(1+X) - X] \right)} \quad (1.4)$$

providing $\eta = \Lambda - \alpha \Lambda^{-1}$. The extra coefficients are introduced to provide

$$Z_{\alpha N}[t.] = Z_N[\eta; \alpha] = Z_N[\Lambda; \alpha] \quad (1.5)$$

to any order of the genus expansion.

A surprising property of the hermitian one-matrix model, which is advocated in Refs.[1, 2], is that it reveals some features of $c = 1$ CFT interacting with 2D gravity. For the case of the Penner model, this property was discovered by Distler and Vafa [4] in the *double-scaling limit* and has been extended to more general models by Tan [5], by Chaudhuri, Dykstra and Lykken [6] and by Gilbert and Perry [7], in particular to the model (1.4) with Λ being proportional to a unit matrix. However, these features of $c = 1$ have been observed in Refs.[1, 2] in genus zero and genus one, identifying α with the cosmological constant, *without* taking the double scaling limit. Therefore, it was conjectured that the hermitian one-matrix model has something to do with the *continuum* $c = 1$ case similar to the Kontsevich model [8] which is associated with the continuum pure 2D gravity ($c = 0$) or to its generalizations [9] which are associated with $c < 1$.

A direct way to verify this conjecture is to study correlation functions of the loop operators. For the hermitian one-matrix model with an arbitrary (not necessarily symmetric) potential, the one-loop correlator has been known in genus zero for a long time [10], while the two- and three-loop correlators were obtained in Ref.[11]. The genus one contribution to the one-loop correlator was explicitly calculated in Ref.[12] for a quartic symmetric potential and in Ref.[1] for an arbitrary symmetric potential. On the other

hand, much is known now about $c = 1$ correlators since pioneering works by Kostov [13] and Boulatov [14]. An incomplete list of references includes [15]–[24].

In the present paper we develop an iterative algorithm for calculating the genus expansion of the hermitian $N \times N$ one-matrix model (= the Penner model in an external field). We introduce the moments, I_p and J_p , of the external field as well as the ‘basis vectors’, $\chi^{(n)}$ and $\psi^{(n)}$, which are determined by a recursion relation and diagonalize the iterative procedure. We prove that the genus g contribution to the partition function depends only on I_p and J_p with $p \leq 3g - 2$ while the genus g contribution to the m -loop correlator depends on I_p and J_p with $p \leq 3g - 2 + m$. We present the explicit calculation of the partition function and the one-loop correlator for an arbitrary potential in genus one. We compare the correlators for the hermitian one-matrix model with those at zero momenta for $c = 1$ CFT and show an agreement of the one-loop correlators (this correlator in $c = 1$ is the only one which vanishes at non-zero momenta) in genus zero. We did not find such an agreement for the two-loop correlator.

2 The iterative scheme

We propose in this section a general iterative procedure for calculating higher genus contributions to the hermitian one-matrix model. We shall be solving, iteratively in $1/N^2$, the loop equation

$$\int_{C_1} \frac{d\omega}{2\pi i} \frac{V'(\omega)}{(\lambda - \omega)} W(\omega) = (W(\lambda))^2 + W(\lambda; \lambda), \quad (2.1)$$

where

$$V(\lambda) = \sum_{k=0}^{\infty} t_k \lambda^k, \quad \frac{\delta}{\delta V(\lambda)} = - \sum_{k=0}^{\infty} \lambda^{-k-1} \frac{\partial}{\partial t_k} \quad (2.2)$$

and

$$W(\lambda) = \frac{\delta}{\delta V(\lambda)} \log Z[t.], \quad W(\lambda; \lambda) = \frac{\delta}{\delta V(\lambda)} W(\lambda) \quad (2.3)$$

with the partition function $Z[t.]$ given by Eq.(1.1). The one-loop correlator $W(\lambda)$ obeys the normalization condition

$$\lambda W(\lambda) \rightarrow \alpha N \quad \text{as} \quad \lambda \rightarrow \infty. \quad (2.4)$$

While we consider in this paper the loop equation (2.1), it should be noticed that a similar iterative procedure can be formulated for the equivalent Schwinger–Dyson equation

$$\left\{ \frac{\partial^2}{\partial \eta_i^2} + \sum_{j \neq i} \frac{1}{\eta_i - \eta_j} \left(\frac{\partial}{\partial \eta_i} - \frac{\partial}{\partial \eta_j} \right) + N \eta_i \frac{\partial}{\partial \eta_i} - \alpha N^2 \right\} Z[\eta.; \alpha] = 0 \quad (2.5)$$

where the partition function $Z[\eta.; \alpha]$ is given by Eq.(1.2) and $\{t_k\}$ are related to η_i — the eigenvalues of η — by Eq.(1.3). The genus one solution of Ref.[1] was obtained for a symmetric distribution of the eigenvalues by solving Eq.(2.5).

Our goal is to solve Eq.(2.1) iteratively order by order in $1/N^2$ starting with the one-cut genus zero solution [10]

$$W_0(\lambda) = \int_{C_1} \frac{d\omega}{4\pi i} \frac{V'(\omega)}{(\lambda - \omega)} \frac{\sqrt{\lambda^2 + b\lambda + c}}{\sqrt{\omega^2 + b\omega + c}}, \quad (2.6)$$

with b and c given by

$$\int_{C_1} \frac{d\omega}{2\pi i} \frac{V'(\omega)}{\sqrt{\omega^2 + b\omega + c}} = 0, \quad \int_{C_1} \frac{d\omega}{2\pi i} \frac{\omega V'(\omega)}{\sqrt{\omega^2 + b\omega + c}} = 2\alpha N. \quad (2.7)$$

In this way we will obtain the genus expansion of the (logarithm of the) partition function

$$\log Z[t.] = \sum_{g=0}^{\infty} F_g[t.] \quad (2.8)$$

and that of the associated multi-loop correlators.

Following Ref.[1], we introduce the new variables

$$\begin{aligned} I_p &= \int_{C_1} \frac{d\omega}{2\pi i} \frac{\omega V'(\omega)}{(\omega^2 + b\omega + c)^{p+\frac{1}{2}}} - (2\alpha + 1)N\delta_{p0}, \\ J_p &= \int_{C_1} \frac{d\omega}{2\pi i} \frac{V'(\omega)}{(\omega^2 + b\omega + c)^{p+\frac{1}{2}}}. \end{aligned} \quad (2.9)$$

Their relation to the moments of the external field η , which is described by the density of eigenvalues $\rho(x)$, can easily be established by substituting

$$\frac{1}{N}V'(\lambda) = \oint dx \frac{\rho(x)}{x - \lambda} - \lambda \quad (2.10)$$

into Eq.(2.9):

$$\begin{aligned} \frac{1}{N}I_p &= \int dx \frac{x\rho(x)}{(x^2 + bx + c)^{p+\frac{1}{2}}} - \left(\frac{3}{8}b^2 - \frac{1}{2}c\right)\delta_{p0} - \delta_{p1}, \\ \frac{1}{N}J_p &= \int dx \frac{\rho(x)}{(x^2 + bx + c)^{p+\frac{1}{2}}} - \frac{1}{2}b\delta_{p0}. \end{aligned} \quad (2.11)$$

The terms with δ_{p0} and δ_{p1} come from the second term on the r.h.s. of Eq.(2.10) taking the residual at infinity. It is easy to see, expanding the integrands in Eq.(2.11) in $1/x$ that I_p depends on t_k with $k \geq 2p$ while J_p depends on t_k with $k \geq 2p + 1$.

As has been proposed in Ref.[1], F_g depends at $1 \leq g < \infty$ only on I_p and J_p for $p \leq P_g - 1$ (where $P_g = 3g - 1$ as is proven below). This is in contrast to the t -dependence of F_g which always depends on the whole set $\{t_k\}$.

For an iterative solution of the loop equation, it is convenient to introduce the operator $\hat{\mathcal{K}}$ by⁴

$$\hat{\mathcal{K}}\Phi(\lambda) = \int_{C_1} \frac{d\omega}{2\pi i} \frac{V'(\omega)}{(\lambda - \omega)} \Phi(\omega). \quad (2.12)$$

⁴Inserting here Eq.(2.10), one finds the relation with the operator $\mathcal{K}$ from Ref.[1]: $-\frac{1}{N}\hat{\mathcal{K}} = \mathcal{K} + \lambda$, when applied to a function $\Phi(\lambda)$ which decays at infinity faster than $1/\lambda$ (if $\Phi(\lambda) \sim 1/\lambda$ as $\lambda \rightarrow \infty$, -1 should be added to the r.h.s. which comes from the residual at infinity).

It is easy to calculate of how the operator $\hat{\mathcal{K}}$ acts on the set of functions

$$\Phi^{(n)}(\lambda) = \frac{1}{(\lambda^2 + b\lambda + c)^{n+\frac{1}{2}}} \quad (2.13)$$

as well as on $\lambda\Phi^{(n)}(\lambda)$. Let us start with $n = 0$. Comparing with Eq.(2.6), one gets immediately

$$\hat{\mathcal{K}}\Phi^{(0)}(\lambda) = 2W_0(\lambda)\Phi^{(0)}(\lambda). \quad (2.14)$$

To calculate of how $\hat{\mathcal{K}}$ acts on $\Phi^{(n)}(\lambda)$ with $n \geq 1$, let us insert the following expansion of the denominator

$$\frac{1}{\lambda - \omega} = (\lambda + \omega + b) \sum_{k=1}^n \frac{(\omega^2 + b\omega + c)^{k-1}}{(\lambda^2 + b\lambda + c)^k} + \frac{(\omega^2 + b\omega + c)^n}{(\lambda^2 + b\lambda + c)^n} \frac{1}{\lambda - \omega} \quad \text{for } n \geq 1. \quad (2.15)$$

Using Eq.(2.9) we get finally

$$\hat{\mathcal{K}}\Phi^{(n)}(\lambda) = \int_{C_1} \frac{d\omega}{2\pi i} \frac{V'(\omega)}{(\lambda - \omega)} \Phi^{(n)}(\omega) = \sum_{k=1}^n \frac{I_{n+1-k} + (\lambda + b)J_{n+1-k}}{(\lambda^2 + b\lambda + c)^k} + 2W_0(\lambda)\Phi^{(n)}(\lambda). \quad (2.16)$$

Quite similarly, one calculates of how $\hat{\mathcal{K}}$ acts on $\lambda\Phi^{(n)}(\lambda)$:

$$\begin{aligned} \hat{\mathcal{K}}\lambda\Phi^{(n)}(\lambda) &= -J_n + \lambda\hat{\mathcal{K}}\Phi^{(n)}(\lambda) = \\ &= \sum_{k=1}^n \frac{\lambda I_{n+1-k} - cJ_{n+1-k}}{(\lambda^2 + b\lambda + c)^k} + \sum_{k=1}^{n-1} \frac{J_{n-k}}{(\lambda^2 + b\lambda + c)^k} + 2W_0(\lambda)\lambda\Phi^{(n)}(\lambda). \end{aligned} \quad (2.17)$$

Some comments concerning these formulas are in order. The expansion is in integer powers of $1/(\lambda^2 + b\lambda + c)$ due to Eq.(2.15). The sum over k is bounded from above by n by construction. For the symmetric potential $V(\lambda) = V(-\lambda)$, which corresponds to a symmetric $\rho(\lambda) = \rho(-\lambda)$ in Eq.(2.10), when $b = 0$ and $J_p = 0$, Eq.(2.16) recovers Eq.(5.13) of Ref.[1].

Let us now turn to the iterative solution of Eq.(2.1). Substituting the genus expansion for $W(\lambda)$ and $W(\lambda; \lambda)$:

$$W(\lambda) = \sum_{g=0}^{\infty} W_g(\lambda); \quad W(\lambda; \lambda) = \sum_{g=0}^{\infty} W_g(\lambda; \lambda), \quad (2.18)$$

and comparing the terms of the same order in $1/N^2$, one rewrites Eq.(2.1) as

$$(\hat{\mathcal{K}} - 2W_0(\lambda)) W_g(\lambda) = \sum_{g'=1}^{g-1} W_{g'}(\lambda) W_{g-g'}(\lambda) + W_{g-1}(\lambda; \lambda) \quad (2.19)$$

where $g \geq 1$. This equation allows us to calculate $W(\lambda)$ order by order of genus expansion.

As is proven below by induction, the r.h.s. of Eq.(2.19), which is known from previous orders of genus expansion, has a structure of the sum over n up to

$$n_g = 3g - 1 \quad (2.20)$$

of the terms of the type $A_g^{(n)}/(\lambda^2 + b\lambda + c)^n$ and $D_g^{(n)}\lambda/(\lambda^2 + b\lambda + c)^n$ with the coefficients $A_g^{(n)}$ and $D_g^{(n)}$ being some functions of the moments I_p and J_p to be discussed latter. For this reason, it is convenient to introduce two sets of ‘basis vectors’ $\chi^{(n)}(\lambda)$ and $\psi^{(n)}(\lambda)$ which obey

$$(\hat{\mathcal{K}} - 2W_0(\lambda))\chi^{(n)}(\lambda) = \frac{1}{(\lambda^2 + b\lambda + c)^n}, \quad (\hat{\mathcal{K}} - 2W_0(\lambda))\psi^{(n)}(\lambda) = \frac{\lambda}{(\lambda^2 + b\lambda + c)^n}. \quad (2.21)$$

This idea is analogous to the one suggested by Gross and Newman [25] for the unitary matrix model and for the hermitian model with a cubic potential.

We can derive now recursion relations which defines $\chi^{(n)}(\lambda)$ and $\psi^{(n)}(\lambda)$ explicitly. Using Eqs.(2.16) and (2.17), we get

$$\begin{aligned} \chi^{(n)}(\lambda) &= \frac{I_1 - \lambda J_1}{\Delta} \Phi^{(n)}(\lambda) - \\ &\quad - \frac{1}{\Delta} \sum_{k=1}^{n-1} \chi^{(k)}(\lambda) \{I_1[I_{n+1-k} + bJ_{n+1-k}] + J_1[cJ_{n+1-k} - J_{n-k}]\} - \\ &\quad - \frac{1}{\Delta} \sum_{k=1}^{n-1} \psi^{(k)}(\lambda) \{I_1 J_{n+1-k} - J_1 I_{n+1-k}\} \end{aligned} \quad (2.22)$$

and

$$\begin{aligned} \psi^{(n)}(\lambda) &= \frac{\lambda(I_1 + bJ_1) + cJ_1}{\Delta} \Phi^{(n)}(\lambda) - \\ &\quad - \frac{1}{\Delta} \sum_{k=1}^{n-1} \chi^{(k)}(\lambda) \{cJ_1[I_{n+1-k} + bJ_{n+1-k}] + (I_1 + bJ_1)[J_{n-k} - cJ_{n+1-k}]\} - \\ &\quad - \frac{1}{\Delta} \sum_{k=1}^{n-1} \psi^{(k)}(\lambda) \{cJ_1 J_{n+1-k} + (I_1 + bJ_1)I_{n+1-k}\} \end{aligned} \quad (2.23)$$

where

$$\Delta = I_1^2 + bI_1 J_1 + cJ_1^2. \quad (2.24)$$

Eqs.(2.21) and (2.22), (2.23) allow us to restore $W_g(\lambda)$ provided the r.h.s. of Eq.(2.19) has the advertised form. The result reads

$$W_g(\lambda) = \sum_{n=1}^{n_g} \left[A_g^{(n)} \chi^{(n)}(\lambda) + D_g^{(n)} \lambda \psi^{(n)}(\lambda) \right]. \quad (2.25)$$

Notice that we do not add the terms with $n = 0$ which are annihilated by the operator $\hat{\mathcal{K}} - 2W_0(\lambda)$. These terms would contradict the boundary condition (2.4). The expression on the r.h.s. of Eq.(2.25) is analytic everywhere in the complex λ -plane except for the cut coinciding with the one of $W_0(\lambda)$. This is in accordance with the different iterative procedure advocated by Migdal [10] and elaborated by David [26].

It remains to prove that the r.h.s. of Eq.(2.19) indeed has such a form. We see from Eq.(2.25) that the first term on the r.h.s. of Eq.(2.19) has exactly this form order by order

in the genus expansion, while the second term needs a more careful analysis since we have to calculate $\delta W_{g-1}(\lambda)/\delta V(\lambda)$. To do this we need some more formulas.

From Eq.(2.9) one gets

$$\begin{aligned}\frac{\partial I_p}{\partial b} &= -(p + \tfrac{1}{2})[bI_{p+1} + cJ_{p+1} + J_p], \quad \frac{\partial I_p}{\partial c} = -(p + \tfrac{1}{2})I_{p+1}; \\ \frac{\partial J_p}{\partial b} &= -(p + \tfrac{1}{2})I_{p+1}, \quad \frac{\partial J_p}{\partial c} = -(p + \tfrac{1}{2})J_{p+1}\end{aligned}\tag{2.26}$$

while Eq.(2.7), which determines b and c , can be rewritten as

$$I_0 = -N, \quad J_0 = 0.\tag{2.27}$$

One needs as well the following formulas for calculating $\delta/\delta V(\lambda)$:

$$\begin{aligned}\frac{\delta b}{\delta V(\lambda)} &= \frac{\partial}{\partial \lambda} \frac{2}{\sqrt{\lambda^2 + b\lambda + c}} \frac{I_1 - \lambda J_1}{\Delta}, \\ \frac{\delta c}{\delta V(\lambda)} &= \frac{\partial}{\partial \lambda} \frac{2}{\sqrt{\lambda^2 + b\lambda + c}} \frac{(\lambda + b)I_1 + cJ_1}{\Delta},\end{aligned}\tag{2.28}$$

with Δ given by Eq.(2.24), and

$$\begin{aligned}\frac{\delta I_p}{\delta V(\lambda)} &= \frac{\partial}{\partial \lambda} \lambda \Phi^{(p)}(\lambda) - \frac{\delta b}{\delta V(\lambda)} (p + \tfrac{1}{2})[bI_{p+1} + cJ_{p+1} + J_p] - \frac{\delta c}{\delta V(\lambda)} (p + \tfrac{1}{2})I_{p+1}, \\ \frac{\delta J_p}{\delta V(\lambda)} &= \frac{\partial}{\partial \lambda} \Phi^{(p)}(\lambda) - \frac{\delta b}{\delta V(\lambda)} (p + \tfrac{1}{2})I_{p+1} - \frac{\delta c}{\delta V(\lambda)} (p + \tfrac{1}{2})J_{p+1}.\end{aligned}\tag{2.29}$$

These formulas⁵ can be obtained from Eqs.(2.9) and (2.27) using Eq.(2.26).

It is easy to see that the result of acting of $\delta/\delta V(\lambda)$ on W_{g-1} given by Eq.(2.25) has exactly the form discussed above. Moreover, we have presented an explicit algorithm for calculating $W_g(\lambda)$, say by symbolic computer calculations with the only input parameter being g . The explicit results for genus one are presented in the next section.

We perform now a power-counting analysis to express n_g and P_g , which are introduced above, via g . Let us note first that the highest moments I_{P_g} and J_{P_g} emerge on the r.h.s. of Eq.(2.25) from the highest term $1/(\lambda^2 + b\lambda + c)^{n_g}$ on the r.h.s. of Eq.(2.19) according to Eqs.(2.22), (2.23) (they are associated with the $k = 1$ terms). Therefore, one gets $P_g = n_g$. To pass to the next order of the genus expansion, one analyze the structure of the r.h.s. of Eq.(2.19). It is easy to estimate the highest power of each term $W_{g'}(\lambda)W_{g-g'}(\lambda)$ for the solution (2.25), which is known from the previous order, to be $n_{g'} + n_{g-g'} + 1$ while the two-loop correlator gives the power $P_g + 3$ by virtue of Eqs.(2.28) and (2.29). Therefore, one gets $n_{g+1} = n_g + 3$ and finally

$$P_g = 3g - 1\tag{2.30}$$

⁵We have not calculated explicitly $\partial/\partial \lambda$ in order that $\delta/\delta \rho(\lambda)$, which is related to $\delta/\delta V(\lambda)$ by

$$\frac{\delta}{\delta V(\lambda)} = \frac{1}{N} \frac{\partial}{\partial \lambda} \frac{\delta}{\delta \rho(\lambda)},$$

could be extracted easily as well.

since $n_1 = 2$. Notice that both types of terms contribute to the maximal power since $n_{g'} + n_{g-g'} + 1 = 3g - 1$.

According to Eqs.(2.8), (2.3) and (2.28), (2.29), the highest moments which contribute to F_g are I_{3g-2} and J_{3g-2} . An analogous result for the Kontsevich model has been obtained by Itzykson and Zuber [27].⁶ Applying Eqs.(2.28), (2.29) m times to obtain the m -loop correlator, one finds highest moments to be I_{3g-2+m} and J_{3g-2+m} . Thus, we have proven the following

Theorem *The genus g contribution to the partition function depends on I_p and J_p with $p \leq 3g - 2$. The genus g contribution to the m -loop correlator depends on I_p and J_p with $p \leq 3g - 2 + m$.*

3 The partition function and correlators in genus one

We present in this section the explicit results for the genus one one-loop correlator $W_1(\lambda)$ and partition function F_1 for the case of an arbitrary potential $V(\lambda)$ which are obtained according to the algorithm of the previous section.

In genus one there is no sum on the r.h.s. of Eq.(2.19) while [11]

$$W_0(\lambda; \lambda) = \frac{b^2 - 4c}{16(\lambda^2 + b\lambda + c)^2}, \quad W_0(\lambda; \mu) = \frac{1}{2(\lambda - \mu)^2} \left[\frac{\lambda\mu - \frac{1}{2}b(\lambda + \mu) + c}{\sqrt{\lambda^2 + b\lambda + c} \sqrt{\mu^2 + b\mu + c}} - 1 \right]. \quad (3.1)$$

Eq.(2.21) yields immediately

$$\begin{aligned} W_1(\lambda) = \frac{b^2 - 4c}{16} \chi^{(2)}(\lambda) = \frac{b^2 - 4c}{16} \left\{ \frac{\Phi^{(2)}(\lambda)}{\Delta} [I_1 - \lambda J_1] - \right. \\ \left. - \frac{\Phi^{(1)}(\lambda)}{\Delta^2} [-J_1^2 I_1 + I_1^2 I_2 + b I_1^2 J_2 - c J_1^2 I_2 + 2c I_1 J_1 J_2] - \right. \\ \left. - \frac{\lambda \Phi^{(1)}(\lambda)}{\Delta^2} [J_1^3 + I_1^2 J_2 - 2I_1 J_1 I_2 - b J_1^2 I_2 - c J_1^2 J_2] \right\} \end{aligned} \quad (3.2)$$

where we have substituted the explicit form of $\chi^{(2)}(\lambda)$ given by Eqs(2.22) and (2.23). For the symmetric case when $b = 0$ and $J_p = 0$, Eq.(3.2) recovers Eq.(5.18) of Ref.[1].

We are now in a position to integrate Eq.(3.2) (w.r.t. ρ) and find the functional F_1 whose variation generates this $W_1(\lambda)$. First, we integrate over λ using the formulas

⁶It would be suggestive to relate these numbers to the (complex) dimension of the moduli space $\mathcal{M}_{g,m}$, which equals $3g - 3 + m$.

$$\begin{aligned}
\frac{1}{4} \int \frac{dx}{(x^2 + bx + c)^{3/2}} &= -\frac{x + \frac{1}{2}b}{(b^2 - 4c)\sqrt{x^2 + bx + c}} + \text{const}, \\
\frac{1}{4} \int \frac{dx}{(x^2 + bx + c)^{5/2}} &= -\frac{x + \frac{1}{2}b}{3(b^2 - 4c)(x^2 + bx + c)^{3/2}} + \\
&\quad + \frac{8(x + \frac{1}{2}b)}{3(b^2 - 4c)^2\sqrt{x^2 + bx + c}} + \text{const}. \tag{3.3}
\end{aligned}$$

It is worth to note the appearance of the factor $b^2 - 4c$ in the denominators.

Using Eqs.(2.29) and (2.28), we can now integrate $W_1(\lambda)$ explicitly. Namely, it is more or less clear that one can produce coefficients like in Eq.(3.2) by differentiating terms like $\log \Delta$ (the terms with Δ^2 in the denominators originate from the variations (2.29), (2.28)). Moreover, due to the appearance of $\frac{1}{\Delta(b^2 - 4c)}$ term after the integration of $W_1(\lambda)$ over λ , it is reasonable to assume the presence of $\log(4c - b^2)$ term in F_1 . After all these preliminaries, a straightforward, though lengthy, calculation gives the answer

$$F_1 = -\frac{1}{12} \log(4c - b^2) - \frac{1}{24} \log \Delta + \text{const}, \tag{3.4}$$

which, in particular, reproduces Eq.(5.18) of Ref.[1] for the reduced model (the symmetric potential).

4 Comparison with $c = 1$ correlators

We compare in this section the explicit formulas for the loop correlators in the hermitian one-matrix model with those in $c = 1$ CFT. A problem immediately arises that the loop operator in $c = 1$ depends both on the length of the loop l and on momentum p which is associated with a matter field. Since in the hermitian one-matrix model there is no dependence on p , we should put p equal to some value (or sum up somehow over special values of p). The procedure is, however, unambiguous for the one-loop correlator which is non-vanishing only at $p = 0$ due to the momentum conservation. Therefore, our idea is to compare first the one-loop correlators.

The genus zero one-loop correlator, given by Eqs.(2.6) and (2.7), is greatly simplified after differentiating w.r.t. the cosmological constant α :

$$\frac{1}{N} \frac{dW_0(\lambda)}{d\alpha} = \frac{1}{\sqrt{\lambda^2 + b\lambda + c}}. \tag{4.1}$$

Let us put for simplicity $b = 0$ keeping in mind that the general case can be restored by the shift $\lambda \rightarrow \lambda + b/2$; $c \rightarrow c - b^2/4$. The equation (2.7) which determines c can be written in the form of the genus zero string equation

$$\sum_m m t_{2m} \frac{(2m-1)!!}{(2m)!!} (-c)^m = \alpha. \tag{4.2}$$

If we restrict ourselves to the simplest case $t_2 \neq 0$, $t_{2m} = 0$ for $m \geq 2$, Eq.(4.2) gives $c = -\alpha/t_2$ and $c > 0$ corresponds to the ‘upside-down’ potential which is familiar from

the quantum mechanical description of $c=1$ correlators. Moreover, the r.h.s. of Eq.(4.2) is as well the known expression for the (diagonal) resolvent of the Schrödinger operator with the potential $c(\alpha)$ in genus zero. Therefore, we conclude that the genus zero one-loop correlators coincide.

The genus zero two-loop correlator in $c=1$ theory at zero momentum can easily be extracted, applying the Gegenbauer's addition theorem, from the results by Moore and Seiberg [22]⁷:

$$W_0(l_1, p = 0; l_2, p = 0) = K_0\left(\sqrt{c}(l_1 + l_2)\right) \quad (4.3)$$

where K_0 is the modified Bessel function of the third kind.

It is easy to see that the Laplace transform of Eq.(4.3) does not coincide with Eq.(3.1). A possible way out (a pure speculative one) might be to sum up the general formula of Moore and Seiberg [22] over the momenta which corresponds to the discrete special states of $c = 1$ CFT [18, 28, 29]. This problem deserves further investigation.

Acknowledgments

We thank C.Kristjansen for discussions. Yu.M. thanks the NBI high energy theory group for the hospitality at Copenhagen.

References

- [1] L.Chekhov and Yu.Makeenko, *A hint on the external field problem for matrix models*, preprint NBI-HE-92-06 (January, 1992), *Phys.Lett.* **B** in print.
- [2] L.Chekhov and Yu.Makeenko, *The multicritical Kontsevich–Penner model*, preprint NBI-HE-92-03 (January, 1992).
- [3] R.C.Penner, *Bull.Am.Math.Soc.*, **15** (1986) 73; *J.Diff.Geom.*, **27** (1988) 35.
- [4] J.Distler and C.Vafa, *Mod.Phys.Lett.*, **A6** (1991) 259.
- [5] C.-I Tan, *Mod.Phys.Lett.*, **A6** (1991) 1373; *Generalized Penner models and multi-critical behavior*, preprint BROWN-HET-810 (May, 1991).
- [6] S.Chaudhuri, H.Dykstra, and J.Lykken, *Mod.Phys.Lett.*, **A6** (1991) 1665.
- [7] G.Gilbert and M.Perry, *Nucl.Phys.*, **B364** (1991) 734.
- [8] M.L.Kontsevich, *Funk.Anal.&Prilozh.*, **25** (1991) 50 (in Russian).
- [9] S.Kharchev, A.Marshakov, A.Mironov, A.Morozov, and A.Zabrodin, *Unification of all string models with $c < 1$; Towards unified theory of 2d gravity*, preprints FIAN/ITEP (October, 1991).
- [10] A.A.Migdal, *Phys.Rep.*, **102** (1983) 199.
- [11] J.Ambjørn, J.Jurkiewicz, and Yu.Makeenko, *Phys.Lett.*, **251B** (1990) 517.

⁷We thank G.Moore for a e-mail correspondence concerning this formula

- [12] J.Ambjørn and Yu.Makeenko, *Mod.Phys.Lett.*, **A5** (1990) 1753.
- [13] I.K.Kostov, *Phys.Lett.*, **215B** (1988) 499.
- [14] D.V.Boulatov, *Phys.Lett.*, **237B** (1990) 202.
- [15] G.Parisi, *Phys.Lett.*, **238B** (1990) 209.
- [16] D.Gross and I.Klebanov, *Nucl.Phys.*, **B344** (1990) 475.
- [17] S.Ben-Menahem, *Nucl.Phys.*, **B364** (1991) 681.
- [18] D.Gross, I.Klebanov, and M.Newman, *Nucl.Phys.*, **B350** (1991) 621;
U.Danielsson and D.Gross, *Nucl.Phys.*, **B366** (1991) 27;
U.Danielsson, *Symmetries and special states in two dimensional string theory*, preprint PUPT-1301 (December, 1991).
- [19] I.Klebanov and D.Lowe, *Nucl.Phys.*, **B363** (1991) 543.
- [20] S.Das and A.Jevicki, *Mod.Phys.Lett.*, **A5** (1990) 1639.
- [21] K.Demeterfi, A.Jevicki, and J.Rodrigues, *Nucl.Phys.*, **B362** (1991) 173; *Nucl.Phys.*, **B362** (1991) 499; *Mod.Phys.Lett.*, **A6** (1991) 3199.
- [22] G.Moore, *Nucl.Phys.*, **B368** (1992) 557;
G.Moore, N.Seiberg, and M.Staudacher, *Nucl.Phys.*, **B362** (1991) 665;
E.Martinec, G.Moore, and N.Seiberg, *Phys.Lett.*, **263B** (1991) 190;
G.Moore and N.Seiberg, *From loops to fields in 2d quantum gravity*, preprint RU-91-29, YCTP-P19-91 (July, 1991).
- [23] I.K.Kostov, *Phys.Lett.*, **266B** (1991) 42; *Phys.Lett.*, **266B** (1991) 317; *Strings with discrete target space*, preprint SPhT/91-142 (September, 1991).
- [24] G.Mandal, A.Sengupta, and S.Wadia, *Mod.Phys.Lett.*, **A6** (1991) 1465;
S.Das, A.Dhar, G.Mandal, and S.Wadia, *Bosonization of nonrelativistic fermions and W-infinity algebra*, preprint IASSNS-HEP-91/72 (November, 1991); *W-infinity ward identities and correlation functions in the $c = 1$ matrix model*, preprint IASSNS-HEP-91/79 (December, 1991).
- [25] D.J.Gross and M.J.Newman, *Phys.Lett.*, **266B** (1991) 291.
- [26] F.David, *Mod.Phys.Lett.*, **A5** (1990) 1019.
- [27] C.Itzykson and J.-B.Zuber, *Combinatorics of the modular group II. The Kontsevich integrals*, preprint SPhT/92-001 (January, 1992).
- [28] E.Witten, *Ground ring of two dimensional string theory*, preprint IASSNS-HEP-91/51 (August, 1991).
- [29] I.R.Klebanov and A.M.Polyakov, *Mod.Phys.Lett.*, **A6** (1991) 3273;
I.R.Klebanov, *Ward identities in two-dimensional string theory*, preprint PUPT-1302 (December, 1991).